31

33

34

35

36

37

38

39

40

41

42

43

44

45

SEDGEO-03774; No of Pages 15



1

2

3 4

6

8

10

11

12 13

14

15

16

17 18

19

21

22 23

24

25

26

27

28

29

Available online at www.sciencedirect.com



Sedimentary Geology xx (2007) xxx-xxx

Sedimentary Geology

www.elsevier.com/locate/sedgeo

Lagoon subsidence and tsunami on the West Coast of New Zealand

S.L. Nichol ^{a,*}, J.R. Goff ^{b,1}, R.J.N. Devoy ^{c,2}, C. Chagué-Goff ^{b,1}, B. Hayward ^d, I. James ^e

^a School of Geography and Environmental Science, The University of Auckland, Private Bag 92019, Auckland, New Zealand
^b National Institute of Water and Atmospheric Research Ltd., PO Box 8602, Christchurch, New Zealand
^c Department of Geography, University College Cork, Cork, Ireland
^d Geomarine Research, 49 Swainston Road, St. John's, Auckland, New Zealand
^e Private Bag 777, Hokitika, New Zealand

Abstract

Sediment core and trench data from a coastal lagoon on the West Coast of the South Island, New Zealand are used to investigate evidence for co-seismic subsidence and associated tsunami inundation. Physical data are used to document a salt marsh soil buried ~80 cm below the modern sediment surface that is locally covered by a gravelly sand bed. The sediment record also contains geochemical and biological (diatom and foram) evidence for abrupt changes in salinity of lagoon waters that link to subsidence, tsunami flooding and to the open versus closed state of the lagoon tidal entrance. At the local scale, these relationships allow for separation of tsunami evidence from other agents of environmental change in the lagoon. We also propose a conceptual connection between these local changes and regional drivers of landscape development, most notably major earthquakes and resultant pulses in sediment supply to the coast.

© 2007 Published by Elsevier B.V.

20

Keywords: Lagoon; Sediments; Holocene; Earthquake; Subsidence; Geochemistry; Diatom; Foraminifera

1. Introduction

Multiproxy analysis of coastal wetland sediments in the Northern Hemisphere has produced important records of past environmental changes, including extreme events such as earthquakes and tsunamis (e.g., Atwater and Moore, 1992; Shennan et al., 1996). Research has generally focused on sites where a demonstrable linkage mechanism (e.g., Pacific northwest, Atwater, 1987; Clague, 1997; Clague et al., 2000). This work and work on other palaeotsunami deposits elsewhere (Dawson et al., 1988; Bryant et al., 1992; Minoura et al., 1996) have served to considerably improve our geological understanding of palaeoseismic-related signatures preserved in the sedimentary record. In New Zealand, we have the opportunity to test and develop this approach by utilising a largely undisturbed and unstudied coastal sediment record that has formed in a seismically active

can be made between tsunamis and their generating

setting (Goff et al., 2001).

New Zealand sits astride the boundary between two major tectonic plates, the Australian and the Pacific. In historical times seismic activity has ranged from barely detectable earthquakes and tsunamis that have caused no

0037-0738/\$ - see front matter © 2007 Published by Elsevier B.V. doi:10.1016/j.sedgeo.2007.01.019

^{*} Corresponding author. Fax: +64 9 373 7434.

E-mail addresses: s.nichol@auckland.ac.nz (S.L. Nichol),
j.goff@niwa.co.nz (J.R. Goff), r.devoy@ucc.ie (R.J.N. Devoy),
c.chague-goff@niwa.co.nz (C. Chagué-Goff),
b.hayward@geomarine.org.nz (B. Hayward), iljames@minidata.co.nz
(I. James).

¹ Fax: +64 3 348 5548.

² Fax: +353 21 427 1980.

damage, to large-scale earth and sea movements (Eiby, 1982; Hull, 1986). Recent reviews of past tsunamis in New Zealand estimate that large, locally generated events, possibly with a nationwide impact throughout the North and South Islands, occur about once every 500 years (Goff and McFadgen, 2002). The most recent events in New Zealand occurred around the mid 15th century AD and are associated with a cluster of large earthquakes, one of which was an Alpine fault rupture (Goff and McFadgen, 2002).

The Alpine fault is over 400 km long, extending almost the entire length of the South Island (Berryman et al., 1992; Bull, 1996; Norris and Cooper, 1997). This

single right-lateral, oblique slip fault raises the Southern Alps at a rate of about 5–8 m per 1000 years (Bull, 1996) (Fig. 1). The last three major ruptures (magnitudes of ca. $8.0~\rm M_w$) are believed to have occurred in AD 1717 (along a 375 km segment), ca. AD 1630 and ca. AD 1460 (involving a minimum 300 km segment, with a vertical offset of $2.15\pm0.4~\rm m$) (Yetton et al., 1998; Wells et al., 1999). A smaller rupture occurred in AD 1826 (Wells et al., 2001; Cullen et al., 2003).

This study examines the recent sedimentary record of Okarito Lagoon on the West Coast of the South Island, with the aim of testing for evidence of co-seismic subsidence of the lagoon floor and tsunami inundation,




Fig. 1. Location map showing Okarito Lagoon, West Coast of South Island, New Zealand.

given the context of frequent Late Holocene seismicity. To achieve this we undertook a multiproxy analysis of sediment cores and trenches from the lagoon. This paper is an extension to earlier work on the same sample set published by Goff et al. (2004).

2. Okarito Lagoon

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89 90

91

92

93

94 95

96

97

98

99

101

102

103

104

108

109

110

111

112

113

114

115

121

Okarito Lagoon, Westland (43°11'S, 170°14'E) is one of the largest estuarine inlets on the West Coast of the South Island. It is 10 km long and 2-4 km wide with an open water area of ~20 km² and fringing salt marsh area of $\sim 10 \text{ km}^2$ (Fig. 1). The lagoon is separated from the sea by a narrow sand barrier with a maximum elevation of ~4 m. The width of the barrier decreases southward from 500 m to 100 m at Okarito village, where an ephemeral tidal inlet provides the only direct connection to the open coast. From vertical aerial photographs it is evident that the lagoon has a complex history of barrier breaching, with four previous tidal inlets ending as blind channels against the landward side of the barrier (Fig. 2). The form of these channels has been maintained as surrounding marshes have developed upon probable relict flood tidal delta surfaces. Our investigation focused on the southernmost marsh island around which the active inlet channel bifurcates, and on the tidal flat to the south of the main channel. Tidal range on the coast varies from 1 m on neaps to 2.1 m on spring tides. Holocene sea-level history for the West Coast is poorly understood in detail, so we assume a similar history for elsewhere in New Zealand which records stillstand since ca. 6.5 ka (Gibb, 1986).

Rainfall in the region exceeds 10,000 mm/year, resulting in high erosion rates (5-12 mm/year) of the fractured, easily eroded schists and glacially derived materials found to the east of the Alpine fault (Berryman et al., 1992; Norris and Cooper, 1997). To the west of the Alpine fault, the overprint of Late Quaternary glacial activity is evident in the large moraines that surround Okarito Lagoon (Griffiths and McSaveney, 1986). Freshwater flow into the lagoon is derived from three sub-catchments; Okarito Forest (9700 ha) which supplies six small creeks that drain the moraines and enter along the northeast shoreline, and; Lake Mapourika (ca. 9000 ha) and Waitangi-taona River (8300 ha) catchments which drain to Okarito River and Deep Creek at the southern end of the lagoon. These catchments extend further inland to drain terrain formed in schist and glacial outwash and are traversed by the Alpine fault in their upper reaches (Fig. 1). The majority of the sediment load is captured in Lake Mapourika and Lake Wahapo, however, and it is the moraines that




Fig. 2. Aerial photograph mosaic (unrectified) for Okarito Lagoon taken January 1988, showing core and trench sites for this study and interpreted former tidal inlets (marked as blind channels). Source: Land Information New Zealand, courtesy Department of Conservation. Crown Copyright Reserved.

124

125

126

127

128

129

130

131

132

133

134

135

137

138

139

140

141

144

145

147

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

provide the main source of coarse material entering Okarito Lagoon (MacPherson, 1981). At its mouth, Okarito River is constructing a small (~1 km²) delta that has infilled the southeast corner of the lagoon. A raised shoreline bench, estimated at 1 m above present water level, fringes the landward side of Okarito Lagoon and is backed by shallow (<5 m) caves cut into cliffed sections of moraines.

An additional key component to landscape development in the West Coast region is seismicity. In particular, ruptures on the Alpine fault have been shown to have resulted in widespread forest destruction due to ground shaking (Wells et al., 1998, 2001; Cullen et al., 2003), river aggradation (Yetton et al., 1998), and rapid coastal dune building (Goff and McFadgen, 2002); the latter two being direct consequences of increased sediment yield from hillslopes following earthquake-induced landslides. It is in this context of strong seismic overprinting on the coastal landscape that we recognized the potential for a record of seismic events to be recorded in Okarito Lagoon.

3. Data sources and analytical techniques

Sediment cores and trenches were used to investigate the shallow stratigraphy of southern Okarito Lagoon. Cores were collected by the vibracore technique at two sites from the lagoon-marsh interface and close to high water mark (Fig. 2). Compaction was measured prior to core recovery. In the laboratory, cores were split lengthwise, logged and sub-sampled for grain size, organic content, micro- and macrofossils, geochemical, and radiocarbon analyses. Trenches (1-3) were dug by hand to confirm the lateral continuity and character of sedimentary units observed in the cores. Additional sediment samples were collected from trenches. Grain size was measured on 23 core and 12 trench sub-samples using a laser particle sizer (GalaiTM) system that determines particle size using the time-of-transition principle (Molinaroli et al., 2000). Results are reported for particle volume measurement. For trench 1, we determined grain size on two samples by mechanical sieving at half-phi intervals from -1 to 4 phi. Organic content of cores was determined via loss-on-ignition (LOI) treatment of 17 sub-samples by ashing at 500 °C for 4 h. Results are reported on a dry weight basis. Elemental analysis of seven sub-samples from core 6 was undertaken using ICP-AES, following sample preparation by a multiacid 'total' digestion. Geochemical data have been normalised for grain size following Loring (1991).

For diatom analysis, 15 sub-samples from core 6 were prepared following standard techniques (e.g., Bat-

tarbee, 1986). Fossil diatoms were mounted in Naphrax and counted using light microscopy at a magnification of × 1000. A minimum count sum of 600 diatom valves was used. Identification of diatom taxa and their palaeoenvironmental interpretation is based upon established floras and other texts, including Hustedt (1927-1966, 1930, 1957), Van der Werff and Huls (1957-1974), Cleve-Euler (1951-1955), Krammer and Lange-Bertalot (1986-1991), Hendey (1964), Round et al. (1990), Simonsen (1967), De Wolf (1982), Admiraal (1984), Denys (1991a,b), Vos and de Wolf (1993a,b, 1994), Foged (1977, 1978, 1979). John (1983), Crosby and Wood (1958, 1959), Wood et al. (1959), and Wood (1961, 1963). Results are presented as a percentage frequency diagram, showing the key diatom species within salinity (Halobion) groupings.

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

For foraminiferal analysis, eight 10-cm³ samples of sediment were processed from cores 4 and 6. Samples were washed over a 63-µm sieve to remove mud and heavy liquid floatation used to concentrate foram tests in sandy samples. Census counts of all benthic forams were made for each sample. Estimates of the tidal elevation at which each foram species was deposited were made using the Modern Analogue Technique (Hayward et al., 2004), based on relative abundance data for 250 benthic foram fauna from modern estuaries and coastal lagoons around New Zealand (data from Hayward et al., 1999a). Similar techniques to determine estimates of tidal elevation based upon foram assemblages have been used elsewhere (Scott and Medioli, 1986; Van de Plassche et al., 1998). In this study, elevation estimates are presented as tidal range derived from the five most similar modern faunas in the analogue set. The reliability of these estimates depends on a number of factors, including the range of tidal levels; depths and environments represented by the analogue samples; and the breadth of the tidal and depth ranges of the major taxa. Previous studies (e.g., Hayward et al., 1999b) have shown that the most precise tidal ranges can be obtained near high tide level from marsh faunas, with far broader ranges observed in intertidal mud and sand flats and in subtidal environments. Some taxa with restricted high tidal ranges in near normal salinity situations (e.g., Haplophragmoides wilberti, Trochamminita salsa, Miliammina fusca) are known to live in abundance through the entire tidal range and also subtidally in more brackish environments (e.g., Hayward et al., 1999a). Thus an assessment of the setting and probable salinity of the lagoon or estuary at the time is important in estimating tidal level.

Foram data also provided for calculation of an artificial salinity index (SI). This was done using detrended

234

235

236

237

238

239

240

241

242

243

244

and canonical correspondence analysis, based on the relative abundance of foram species following Hayward et al. (2004). SI estimates range between 0 (freshwater) and 10 (normal marine salinity).

228 **4. Results**

229 4.1. Stratigraphy and sediment texture

All five sample sites preserve a similar stratigraphic record within the upper metre (Figs. 3-5). The

232 bulk of the sediment matrix comprises massive beds

of fine to medium sand and silt with local shell and wood fragments. Within this, each core and trench recovered a buried marsh soil at variable depths, ranging from 50 cm (trench 3) to 88 cm (trench 1) with depths in the cores falling within this range. The soil unit is olive-brown to grey-olive silt that encloses fine root hairs and wood fragments. It is in sharp upper contact with overlying sediments that are distinguished by a colour change to light grey and absence of organic material, an observation supported by LOI results which show a 1–2% decrease across this contact (Fig. 6).




Fig. 3. Graphic logs for cores 4 and 6.

S.L. Nichol et al. / Sedimentary Geology xx (2007) xxx-xxx




Fig. 4. Graphic logs for trenches 1-3.

In trench 1, the buried soil is overlain by a 23-cm-thick bed of shell-bearing gravel and coarse sand. Gravels are rounded and shell remains include whole and articulated bivalves near the base of the unit that become more commonly broken toward the upper contact at 65 cm depth. Shell species are dominated by the cockle Austrovenus stutchburyi, with rare examples of the intertidal bivalve Mactra ovata and the gastropod Amphibola crenata. In addition, remains of macro

algae, most likely seaweed, were recovered. On the landward side of the lagoon (core 4 and trench 2) sediment above the soil comprises 20–30-cm-thick beds of very fine sand that are massive and carry shell imprints and wood fragments. In trench 2, the contact is on an irregular surface and grey fine sands have filled vertical cracks that penetrate 10 cm into the buried soil; these cracks may have originated as burrows or, more likely, via desiccation of the soil when exposed subaerially. In

254

255

256

257

258

259

260

261

262

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308




Fig. 5. (A) Core 6 showing soil surface buried 77 cm below modern tidal flat and articulated bivalve (Austrovenus stutchburyi) dated to 900±60 years BP; (B) trench 1 showing buried soil overlain by rounded pebbles, gravel and shell, grading upward to very fine sand. Scale card is 10 cm long; (C) trench 2 showing buried soil overlain by very fine sand; arrow indicates infilled desiccation crack or burrow. Tape is 50 cm long; (D) trench 3 showing buried soil overlain by fine sand.

trench 3, the buried soil is overlain by a 4-cm-thick bed of coarse sand and medium-sized gravel that grades upward to a 40-cm-thick deposit of very fine sand,

266 4.2. Geochemistry

267

268

273

274 275

276

277

278

279

280

281

283

284

285

A suite of elements assayed from core 6 show a clear peak in concentration within the silty sands that overlie the buried soil (Fig. 6). Thus, iron (Fe), sulphur (S), titanium (Ti), strontium (Sr), barium (Ba), calcium (Ca) and sodium (Na) all increase markedly at 77 cm depth and then decrease within the successive bed of finemedium sand to 20 cm depth. Although S occurrence in sediments is mostly related to post-depositional diagenetic processes. Fe-rich sediments are more likely to fix S than Fe-poor sediments. Thus S and Fe can be used together as indicators of palaeosalinity in sediments (Thomas and Varekamp, 1991; Daoust et al., 1996; Chagué-Goff and Goff, 1999). Na has also previously been used as a proxy for marine influence in marsh and lagoon systems, because of the relatively high Na content in seawater compared with freshwater (López-Buendía et al., 1999; Chagué-Goff et al., 2002). Similarly, Sr, Br and Ca also occur in higher concentrations in marsh and wetland sediments inundated by saltwater

(Minoura et al., 1994; Chen et al., 1997). Titanium is used here as an indicator of heavy mineral concentration (e.g., magnetite). Therefore, the peak Ti concentration directly above the buried soil within core 6 is interpreted to record the preferential sorting and concentration of heavy minerals. In a lagoon, this particular response would most likely occur during a high energy event such as initiated by tsunami, storm surge or large river flood. Collectively, the measured elements all indicate a sudden rise in the salinity of lagoon waters, followed by a gradual return to brackish conditions.

4.3. Diatoms

Diatom analysis of core 6 provides further evidence of palaeoenvironmental change for Okarito Lagoon (Fig. 7). Fossil diatoms are generally well preserved and abundant, providing a flora of >240 taxa. In the lower part of the core (87–100 cm depth) brackish water taxa (mesohalobion, M; oligohalobion, OI-indifferent and OH-halophile) comprise >50% of the diatom assemblage. Melosira juergensii and Mastogloia elliptica dominate the M taxa, with lower but significant values of Mastogloia pseudoexigua, Cyclotella operculata var. mesoleia and Rhopolodia gibberula (OH — indicative




Fig. 6. (A) Geochemical data for core 6, also showing trends in silt and organic content. (B) Trends in silt and organic content for core 4.

309 of sandy conditions). The OI group, rich in Cocconeis placentula and Cyclotella stelligera, maintains frequen-310 cies >32% throughout the interval. There is a distinct 311 background marine influence (polyhalobion, P), but at 312 frequencies <15%. Marine diatoms are represented 313 by Diploneis suborbicularis, Raphoneis surirella, 314 Raphoneis minutissima, Thalassiosira eccentrica and Paralia sulcata. These are littoral species, as are many 316 of the M group; together representing shallow marine-317 brackish water of sandy foreshore, sand flat and mud flat environments (Denys, 1985). Open coast marine species are also recorded in the sediments, confirming tidal exchange at the site. These marine diatoms are 321 sparse and often broken, indicating their presence is probably the result of progressive inwashing and reworking of more open water, sandy marine sediments. 324

The interval between 67 and 87 cm depth is distinguished by alternating peaks of marine (P) and brackish-freshwater (OH, OI) groups. Marine diatoms have peak concentration at 85 cm (20%) and 73 cm (27%), the former associated with the buried soil and the latter with the overlying fine sand. Species typical of sandy-mud flat environments are prominent in these apparently marine-enriched sediments, with increased values of Raphoneis spp., Cocconeis scutellum, Nitzschia granulata, N. punctata, Striatella unipunctata and Licmophora spp. Many other marine species occur in low numbers, including Cerataulus turgidus, Actinocyclus octonarius var. crassus, Grammatophora oceanica, Triceratium spp. and Navicula lyra.

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

A decline in marine taxa (P) occurs above 85 cm depth through the buried soil and high values of brackish

S.L. Nichol et al. / Sedimentary Geology xx (2007) xxx-xxx




Fig. 7. Summary plot of diatom salinity (halobion) groups for core 6. Polyhalobion (P) — marine conditions, mesohalobion (M) — full brackish water conditions, oligohalobion (OI) — freshwater, salt indifferent; oligohalobion (oh) — some brackish water tolerance; halophobous (H) — freshwater conditions.

(OH) and freshwater taxa (OI) are recorded, with individual maxima for R. gibberula, C. placentula and Navicula tripunctata. Full brackish water species (M) increase above the soil with Mastogloia spp., Melosira jeurgensii reaching peak frequencies at 75–73 cm, with lower values of Synedra fasciculata and Nitzschia sigma. Many of these OI–M group diatoms are well adapted to intertidal–salt marsh conditions and are tolerant of drying-air exposure.

Between 67 and 30 cm depth, freshwater (OI) species dominate and frequencies of brackish (M) diatoms decline from >30% to $\sim20\%$. Above 67 cm the marine

signal collapses to <2% above 50 cm. The OI and OH diatoms (together >60% at the base of this interval) are joined by a range of other fresh-brackish water species, increasing to near 80% at 30 cm. These include many *Achnanthes* spp., with *Achnanthes hungarica* replacing *C. placentula* as the dominant taxon at the top of the zone. The diatom assemblage suggests an overall progressively shallowing, fresh-brackish water and river flood-dominated environment.

Many of the fresh and brackish water diatom valves (OI/OH and M groups) are relatively unworn and whole valves dominate the general diatom assemblages (Fig. 7).

368

369

370

371

373

377

378

379

380

381

382

383

384

385

386

387

388

389

391

392

393

t1.1

t1.2

t.1.20

t1.21

t1.22

This lack of breakage is unusual given the potential for high-energy conditions and the reworking of sediments at the site. Many of the OI-M diatoms however represent low to medium salinity environments (Cl <20 ppt) and are predominantly of local to *in situ* origin, with growth under intertidal—subaerial conditions. Peaks in key freshwater (OI) species between 87 and 30 cm are associated with terrestrial plant debris most probably introduced during river flooding. Overall, the diatom record from core 6 shows deposition generally in brackish to brack-ish-freshwater environments of estuary-lagoonal—marsh settings. Toward the top of the sedimentary succession a marked decline in the percentages of marine species and the associated increase in freshwater species indicate an increasing isolation from the sea.

The higher resolution diatom analysis between 67 and 87 cm shows at least one distinctive marine 'event' in the sedimentary sequence. It is worth noting that, if a lower sampling resolution had been maintained, then the extremes in diatom values (75-70 cm) would not have been apparent. Two peaks in marine diatoms are recorded. The first, at 85 cm, coincides with the base of the buried soil. In the context of a consistent marine diatom signal this peak is most likely an artifact of the stratigraphy and sedimentary processes. Marine diatoms tend to be more heavily silicified and resistant to destruction. Local reworking of sediments to concentrate the marine diatoms present may be the explanation, as supported by the increase in broken valves at this level. The second marine peak (at 73 cm) is more distinctive and associated with the sand bed overlying the buried

soil. This peak is bracketed by higher concentrations of fresh-brackish water diatoms (e.g., 80 cm within the soil, 71 cm in fine sand) that we interpret to represent the prevailing background shallow water, intertidal to salt marsh environment at the site.

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

4.4. Foraminifera

Forams extracted from cores 4 and 6 are used here to reconstruct palaeo-elevation ranges for depositional surfaces (Table 1). A total of 10 taxa are reported, in varying abundance and diversity through the sediment record. In core 6, a sample taken directly below the contact between the buried soil and overlying sand (79-81 cm) is co-dominated by agglutinated faunas that indicate weakly brackish conditions (Ammotium fragile, H. wilberti) on a lagoon or marsh surface that was close to mean high tide level (range 2-2.2 m above MSL). Within the overlying sand (71-73 cm), the fauna is strongly dominated by calcareous taxa (i.e., Ammonia aoteana) indicative of increased salinity and a broader elevation range spanning subtidal (2.5 m below MSL) to mid-tide (1.75 m above MSL). Four samples taken from the interval 64-5 cm in core 6 all indicate a return to lower salinity, based on consistently high values for H. wilberti and THE strong presence of A. fragile and M. fusca. Estimated elevations range from mid-tide (1.9 m above MSL) to mean spring high tide level (2.2 m above MSL), most likely in a brackish marsh environment.

Foram concentrations from the two samples taken from core 4 are very low, allowing estimation of tidal

Table 1
Foraminifera data for Okarito cores 4 and 6

3		Core 6 ^a ,						Core 4 ^b		
4	Depth (cm)	5-7	50-52	57-59	62-64	71–73	79–81	101-103	54-56	76–78
5	Number of species	5	3	2	5	6	5	5	1	4
6	Total foram count	104	201	3	25	163	360	250	7	11
7	Relative abundance %									
3	Ammobaculites exiguus	0	0	0	0	0	0	0	0	46
9	Ammonia aoteana	0	0	0	20	77	1	35	0	0
10	Ammotium fragile	13	54	0	4	5	34	19	0	18
11	Elphidium excavatum f. clavatum	0	0	0	0	1	0	4	0	0
12	E. excavatum f. excavatum	0	0	0	0	1	0	0	0	0
13	Haplophragmoides wilberti	64	45	67	68	14	30	34	0	18
4	Jadammina macrescens	1	0	0	0	0	0	0	0	0
.5	Miliammina fusca	21	1	33	4	2	14	8	100	18
16	Pseudothurammina limnetis	0	0	0	0	0	21	0	0	0
7	Trochamina inflata	1	0	0	4	0	0	0	0	0
18	Elevation (m above MSL)	1.9-2.2	2.0 - 2.2	n/a c	2.2	-2.5 - 1.75	2.0 - 2.2	-2.5-1.15	n/a	-2.5-1.75
19	Salinity index	2.6	2.4	2.6	3.3	5.5	2.2	6.2	3.2	4.3

^a Core 6 from edge of *Leptocarpus similis* salt marsh at neap high tide level, ~1.8 m above MSL.

b Core 4 from Leptocarpus similis salt marsh at mean high tide level, ~2 m above MSL.

c n/a indicates insufficient foram preservation to allow application of MAT.

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

t2.1 Table 2 t2.2 Radiocarbon results for Okarito cores

t2.3	Laboratory code	Core	Depth (cm)	dC ¹³	C ¹⁴ age years BP	Calibrated age BP (2 sigma)	Material dated		
t2.4	Wk8989	4	65	-26.3 ± 0.2	1800±60	1870-1560	Root from buried soil		
t2.5	Wk8987	6	78	-21.5 ± 0.2	2170 ± 70	2340-1950	Bulk soil/organics — buried soil		
t2.6	Wk8618	6	82	1.1 ± 0.2	$10,580 \pm 60$	12,280-10,880	Articulated shell — Austrovenus stutchburyi		
t2.7	Wk8986	6	82	-0.6 ± 0.2	900±60	630-455	Articulated shell — Austrovenus stutchburyi		

t2.8 Calibrations based on Stuiver et al. (1998) using a ΔR of -30 ± 15 for marine shells (McFadgen and Manning, 1990).

425 level for only one sample (76–78 cm, Table 1). The fauna from this sample indicate a wide elevation range, from 426 subtidal (2.5 m below MSL) to mean high water neap 427 428 (1.75 m above MSL).

In summary, the forams identified from these cores provide evidence for abrupt changes in salinity during the accumulation of the upper metre of Okarito Lagoon infill. We have confidence in this evidence because the preservation state of tests is moderate to good, there is no evidence for reworking of older tests into the sampled sediments and open marine fauna were not recorded; together indicating an intact and in situ faunal assemblage that reliably reflects palaeo-salinity conditions at the sample sites.

439 4.5. Radiocarbon ages

429 430

433

434

435

436

451

456

457

Radiocarbon results for four samples taken from cores 440 4 and 6 are presented in Table 2. Samples of buried soil and root material from within yielded similar uncalibrated ages of 1800±60 years BP (Wk8989; core 4) and 2170±70 years BP (Wk8987, core 6). Two specimens of 444 articulated bivalve, taken from the base of the buried soil 445 in core 6, yielded contrasting ages of 10,580±60 years BP (Wk8618) and 900 ± 60 BP (Wk8986). We view the older age as unreliable on the basis that sea level was about 30 m lower than present at ~ 10 ka and the area now occupied by Okarito Lagoon would have ice-marginal terrain. Contamination of sample Wk8618 by old carbon is therefore assumed and the age rejected. The true 452 Middle Holocene age for the lower part of the lagoon 453 infill is supported by a C¹⁴ age of 6380±60 years BP on 454 articulated cockle (A. stutchburyi) recovered from 326 cm depth in core 6 as part of a previous study (Goff et al., 2004).

5. Interpretation and discussion 458

The preservation of a buried soil at similar depths 459 below the modern salt marsh and fringing tidal flat of Okarito Lagoon is strong evidence for localized sub-461

sidence. Results from diatom- and foram-based reconstruction of palaeo-salinity support an interpretation of abrupt changes in local environmental conditions across the contact between the buried soil and overlying deposits. Thus, we recognize an increase in the salinity of lagoon waters associated with increased water depth of approximately 50 cm. Further evidence for increased tidal incursion at core site 6 is given by geochemical data that record a marked increase in a suite of elements to indicate marine-enriched lagoon waters (especially S, Na, Ca and Ba), associated with sediments directly above the buried soil. We note, however, that these sediments do not contain any open marine foraminifera species, so that evidence for transport of material from the open coast is lacking.

Sediments directly overlying the buried soil range in grain size from coarse sand and gravel (trench 1 and 3) to silty fine sand (core 4 and 6, trench 2). Our sampling does not allow for detailed mapping of these deposits, but local variability of grain size such as between core 6 and trench 1 (a distance of ~ 100 m) suggests that sediment texture may vary with proximity to tidal channels, with channels being the likely source for local reworking of coarse-grained sediments. These deposits grade-upward to silty fine sands that contain an increasingly dominant freshwater diatom assemblage, which suggests that the site became less open to tidal influence following deposition of the coarser bed. Abrupt declines in geochemical indicators of saline conditions point to a similar transition. On this basis, we interpret the finer sediments as tidal flat deposits, formed under regular, low-energy conditions.

Here we propose a scenario to account for the observed changes in the shallow stratigraphy of Okarito Lagoon. Given the proximity of Okarito Lagoon to the Alpine fault, the most likely mechanism for subsidence of the lagoon floor is earthquake activity. Co-seismic subsidence has not previously been recognized for the West Coast, but is well established for similar tectonic settings elsewhere; most notably the Pacific Northwest coast of the USA (Atwater, 1987). Subsidence of coastal

505

506

507

508

509

510

511

517

518

519

520

521

522

523

524

525

528

529

531

532

533

534

535

536

538

539

540

542

543

544

545

546

547

548

549

550

553

554

lowlands has also been associated with tsunami inundation of these coasts, with sedimentary evidence preserved as anomalous deposits (sand sheets) across the subsided surface (Atwater and Moore, 1992).

This leads to the question — did co-seismic subsidence of 0.5 m in Okarito Lagoon trigger a local tsunami that left a sedimentary record? Based on the results presented, we highlight the following as potential tsunami evidence: (1) the gravel and pebble deposits in trench 1, and to a lesser degree, the coarse sand and fine gravel in trench 3 are a compelling indicator of highenergy flow (metres per second) across the subsided soil surface; (2) the abrupt rise in lagoon salinity as recorded by geochemical data, diatoms and forams is consistent with, though not exclusive to, sudden incursion of marine waters, and; (3) the re-establishment of reduced salinity conditions in sediments above the coarsegrained deposits, as indicated by diatoms particularly, strengthens the interpretation that higher salinities are not the norm for Okarito Lagoon.

In evaluating the case for a tsunami in Okarito Lagoon (and adjacent coast), we must also consider other scenarios as explanation for the observed sedimentary record. For these alternatives, we draw upon geomorphic and historic evidence. We noted earlier that the barrier to Okarito Lagoon has a series of blind channels that terminate behind the modern foredune (Fig. 2). These channels are interpreted as former tidal inlets. Given that the current (micro) tidal prism of the lagoon only requires one inlet, we assume that the lagoon has always functioned as a single-inlet system, the location of the inlet has shifted and that at times the lagoon has remained closed. Support for this assumption is found in the raised shoreline bench and shallow caves that fringe the landward side of Okarito Lagoon. The simplest explanation for this raised shoreline is that the lagoon was up to 1 m deeper and closed to tidal exchange. An alternative is that the bench formed during the Middle Holocene sea-level highstand. However, the highstand remains to be convincingly documented for this coast. Moreover, recent dendrochronological work on the rimu (Dacrydium cupressinum) and kahikatea (Dacrycarpus dacrydioides) forest that has colonized the bench indicates that the oldest trees post-date AD 1832 (Goff et al., 2004). Thus, it is likely that lowering of the lagoon water level occurred shortly prior to ca. AD 1830 and was presumably associated with barrier breaching to re-establish tidal exchange, not a change in relative sea level. Goff et al. (2004) propose tsunami as the mechanism for barrier breaching and identify a known tsunami in AD 1826 as the likely event.

Further indirect evidence for subsidence and tsunami along this coast is found in historical accounts from the mid-nineteenth century and is summarized in Goff et al. (2004). Of these accounts, keys for this study are two reports of a drowned Maori village at Poherua (aka Saltwater) Lagoon 11 km to the north of Okarito Lagoon. The first report was from the explorer Thomas Brunner in 1848 (republished in Pascoe, 1952) and the second in 1864 by Dobson, who later wrote, "At Poherua lagoon there had been at one time a Maori village of considerable size, the stumps of the posts of the houses showing plainly...The land had sunk..., as these traces could be seen at dead low water, spring tides." (Dobson, 1930, p.72). Dobson also noted that local Maori were unaware the village had ever existed, which suggests to us that its subsidence and abandonment occurred well before the 19th century. This site was last verified in the 1980's (New Zealand Forest Service, in press).

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

Returning to the point regarding lagoon closure, this will be achieved by accretion of the beach berm on the barrier, leading to blocking of the inlet (a process that occurs about once a decade today). Clearly, this requires a positive sediment budget in the littoral zone and we recognize the large gravel-bed rivers (Waiho River and Cook River) to the south of Okarito Lagoon as obvious long-term sediment sources for barrier construction. However, it is unlikely that this supply has been at a consistent rate. There is good evidence from the Haast coast, ~100 km to the south of Okarito Lagoon, that barrier construction has been episodic and linked to fluctuations in sediment discharge from rivers draining the Southern Alps. Reconstruction of the chronology of dune-ridge formation by Wells and Goff (2006) has identified a link to known earthquakes on the Alpine fault. It follows that these events have led to increased sediment supply to rivers from earthquake-induced landslides, and in turn, to the coast. Given this eventbased fluctuation in sediment supply to barriers, periodic closure of lagoons along the West Coast is an expected outcome.

To summarize, fluctuations in the salinity and water level of Okarito Lagoon can be readily accounted for by periodic opening and closing of the tidal inlet entrance. However, this cannot be used to explain the placement of gravel and coarse sand at relatively distal locations in the lagoon, nor the preservation of a buried salt marsh soil. The shallow depth of the gravel and sand deposits also precludes transport as a lag along a tidal channel because flow velocities in a <1 m deep channel inside a microtidal lagoon would not exceed threshold for gravel transport. We also discount river floods as a transport mechanism on the basis that the competence of the

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699 700

701

702

703

704

607 Okarito River is limited to transport of coarse sand to the delta front, which is on the opposite shore of the lagoon. \sim 1.4 km from the site of trench 1. Tsunami remains the 610 only plausible mechanism for transport of coarse sand and gravel to this relatively quiescent location. And the association with a subsided soil provides the important link to palaeo-seismicity.

614 6. Age of this event?

Establishing the age for a palaeo-tsunami event is 615 616 problematic, with the best results often obtained by dating material in sedimentary units above and below an 617 interpreted tsunami deposit, to "age bracket" the event. 618 619 Radiocarbon ages are often equivocal because of the possible introduction of older reworked carbon (wood and other macro-/microfossil debris) (Dawson, 1994; Goff et al., 1998). We have already discounted one 622 reported age (10,580±60 years BP) on this basis. Acknowledging the further possibility that organic material in the buried soil contains old carbon derived from the 626 catchment, the two radiocarbon ages of 1800±60 years BP and 2170±70 years BP are interpreted as maximum ages for deposition of the enclosing sediments. Our 629 remaining radiocarbon age of 900±60 years BP is de-630 rived from an in situ bivalve preserved within the buried soil. The calibrated age of this sample is in the range 631 630-455 years BP (1320-1495 AD) and represents the 632 633 best age estimate for the subsidence event and tsunami.

This age correlates well with the earliest in a series of 635 recognized ruptures on the Alpine fault, dated to ca. AD 1450 (Bull, 1996; Yetton et al., 1998). Other ruptures are 636 dated to AD 1620, 1717 and 1826, with the latter gene-638 rating a tsunami. These events are also correlated with large-scale forest disturbance due to severe ground shaking and landslides (Wells et al., 1998, 1999, 2001; Cullen et al., 2003). It seems highly likely therefore that the tsunami reported here is linked directly to the mid 15th century event, either as the direct result of subsidence associated with onshore fault rupture, submarine fault rupture (the southern segment of the Alpine fault is submarine) or of submarine landsliding generated by an earthquake. These sources remain to be verified, however. It is also suggested that the mid 15th century event caused sufficient forest destruction to trigger increased sediment supply to the coast and, in turn, to closure of the tidal inlet to Okarito Lagoon.

652 7. Conclusion

634

639

640

642

643 644

645

646

647

648

649

650

653 Multiproxy analysis of a distinct sedimentary succession recovered at five sites in Okarito Lagoon provided 654

for reconstruction of palaeoenvironments that record coseismic subsidence of a salt marsh surface and inundation of that surface by tsunami. Key criteria include a buried soil, associated with major upward changes in grain size, geochemistry, diatom and foraminiferal assemblages. Limited radiocarbon dating of the buried soil suggests that the subsidence occurred in the mid 15th century and was linked to a rupture on the nearby Alpine fault. Of importance to this case is the necessity to separate other possible mechanisms from the subsidence-tsunami interpretation. Thus, geomorphic changes to the barrier inlet are also recognized as an important influence on lagoon water level and salinity, but not as a causal mechanism for the deposits preserved in the lagoon fill. Ultimately, however, there is a link because those changes in barrier form can be potentially traced back to major seismic events in the region. Further work to test this hypothesis will require extension of the record presented here to demonstrate repetition of the process over time and across new study sites.

Acknowledgements

Financial support was provided to this project by the Department of Conservation (Hokitika, Southern Regional Office), West Coast Regional Council, The University of Auckland, the Australian Institute of Nuclear Science and Engineering, the New Zealand Foundation for Research, Science and Technology, and GeoEnvironmental Consultants. We thank Quentin Smith for field work assistance and Anne Hall, Keri Hulme and Debbie McLachlan from Okarito for support with logistics and local information. Bruce McFadgen is thanked for providing calibrations of radiocarbon ages. J.-P. Guilbault, A.R. Nelson, and W.B. Bull provided useful comments on earlier versions of this paper. Many thanks to Dave Tappin for a comprehensive final review that significantly improved the paper.

References

Admiraal, W., 1984. The ecology of estuarine sediment-inhabiting diatoms. Progress in Phycological Research 3, 269-322. Atwater, B.F., 1987. Evidence for great Holocene earthquakes along the outer coast of Washington State. Science 236, 942-944. Atwater, B.F., Moore, A.L., 1992. A tsunami about 1000 years ago in

Puget Sound. Washington. Science 258, 1614-1617. Battarbee, R.W., 1986. Diatom analysis. In: Berglund, B.E. (Ed.), Handbook of Holocene Palaeoecology and Palaeohydrology. Wiley,

Chichester, pp. 527-570. Berryman, K.R., Beanland, S., Cooper, A.F., Cutten, H.N., Norris, R.J., Wood, P.R., 1992. The Alpine fault, New Zealand: variation in Quaternary structural style and geomorphic expression. Annales

Tectonicae VI 126-163

- 705 Bryant, E.A., Young, R.W., Price, D.M., 1992. Evidence of tsunami
 706 sedimentation on the southeastern coast of Australia. Journal of
 707 Geology 100, 753-765.
- 708 Bull, W.B., 1996. Prehistoric earthquakes on the Alpine fault, New Zealand. Journal of Geophysical Research 101 (B3), 6037-6050.
- Chagué-Goff, C., Goff, J.R., 1999. Geochemical and sedimentological
 signature of catastrophic saltwater inundations (tsunami), New
 Zealand. Quaternary Australasia 17, 38–48.
- Chagué-Goff, C., Dawson, S., Goff, J.R., Zachariasen, J., Berryman,
 K.R., Garnett, D.L., Waldron, H.M., Mildenhall, D.C., 2002. A
 tsunami (c. 6300 years BP) and other environmental changes,
 northern Hawke's Bay, New Zealand. Sedimentary Geology 150,
 89–102.
- Chen, Z., Chen, Z., Zhang, W., 1997. Quaternary stratigraphy and
 trace-element indices of the Yangtze Delta, eastern China, with
 special reference to marine transgressions. Quaternary Research
 47, 181–191.
- Clague, J.J., 1997. Evidence for large earthquakes at the Cascadia
 subduction zone. Reviews of Geophysics 35, 439-460.
- Clague, J.J., Bobrowsky, P.T., Hutchinson, I., 2000. A review of
 geological records of large tsunamis at Vancouver Island, British
 Columbia. Quaternary Science Reviews 19, 849–863.
- 727 Cleve-Euler, A., 1951–1955. Die Diatomeen von Schweden und 728 Finnland. Kungliga Svenska Vetenskaps-akademiens Handlingar 729 Serie 4, 2 (1), 3–163 (1951); 3 (3), 3–153 (1952); 4 (1), 3–158 730 (1953); 4 (3), 3–225 (1953); 5 (4), 3–231 (1955).
- Crosby, L.H., Wood, E.J.F., 1958. Studies on Australian and New
 Zealand diatoms, I: planktonic and allied species. Transactions of
 the Royal Society of New Zealand 85, 483-530.
- Crosby, L.H., Wood, E.J.F., 1959. Studies on Australian and New
 Zealand diatoms, II: normally epontic and benthic genera. Transactions of the Royal Society of New Zealand 86, 1-58.
- 737 Cullen, L.E., Duncan, R.P., Wells, A., Stewart, G.H., 2003. Floodplain
 738 and regional scale variation in earthquake effects on forests,
 739 Westland, New Zealand. Journal of the Royal Society of New
 740 Zealand 33, 693-701.
- Daoust, R.J., Moore, T.R., Chmura, G.L., Magenheimer, J.F., 1996.
 Chemical evidence of environmental changes and anthropogenic influences in a Bay of Fundy saltmarsh. Journal of Coastal Research 12, 520-533.
- 745 Dawson, A., 1994. Geomorphological effects of tsunami run-up and746 backwash. Geomorphology 10, 83-94.
- Dawson, A., Long, D., Smith, D.E., 1988. The Storegga slides: evidence from eastern Scotland for a possible tsunami. Marine Geology 82, 271-276.
- Denys, L., 1985. Diatom analysis of an Atlantic-SubBoreal core from
 Slijpe (Western Belgian coastal plain). Review of Palaeobotany
 and Palynology 46, 33-53.
- Denys, L., 1991a. A checklist of the diatoms in the Holocene deposits
 of the western Belgian coastal plain with a survey of their apparent ecological requirements.
 Introduction, ecological code and
 Complete list. Belgische Geologische Dienst, Professional Paper
 1991/2, 246.
- Denys, L., 1991b. A checklist of the diatoms in the Holocene deposits of the western Belgian coastal plain with a survey of their apparent ecological requirements. II. Centrales. Belgische Geologische Dienst, Professional Paper 1991/3, 247.
- 762 De Wolf, H., 1982. Method of coding of ecological data from diatoms
 763 for computer utilisation. Mededelingen Rijks Geologiische Dienst
 764 36, 95-98.
- Dobson, A.D., 1930. Reminiscences of Arthur Dudley Dobson. Whit-combe and Tombs, Wellington.

Eiby, G.A., 1982. Two New Zealand tsunamis. Journal of the Royal Society of New Zealand 12, 337-351.

767

768

769

770

771 772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789 790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

- Foged, N., 1977. Freshwater diatoms in Ireland. Bibliotecha Phycologica 34.
- Foged, N., 1978. Diatoms in Eastern Australia. Bibliotecha Phycologica 41.
- Foged, N., 1979. Diatoms in New Zealand, the North Island. Bibliotecha Phycologica 47.
- Gibb, J.G., 1986. A New Zealand regional Holocene eustatic sea-level curve and its application to determination of vertical tectonic movements. Royal Society of New Zealand, Bulletin 24, 377–395.
- Goff, J.R., McFadgen, B.G., 2002. Seismic driving of nationwide changes in geomorphology and prehistoric settlement—a 15th century New Zealand example. Quaternary Science Reviews 21, 2313-2320.
- Goff, J.R., Crozier, M., Sutherland, V., Cochran, U., Shane, P., 1998.
 Possible tsunami deposits of the 1855 earthquake, North Island, New
 Zealand. In: Stewart, I.S., Vita-Finzi, C. (Eds.), Coastal Tectonics.
 Geological Society Special Publication, vol. 133, pp. 353–374.
 London.
- Goff, J.R., Chagué-Goff, C., Nichol, S., 2001. Palaeotsunami deposits: a New Zealand perspective. Sedimentary Geology 143, 1-6.
- Goff, J.R., Wells, A., Chagué-Goff, C., Nichol, S.L., Devoy, R.J.N., 2004. The elusive AD 1826 tsunami, South Westland, New Zealand. New Zealand Geographer 60, 14-25.
- Griffiths, G.A., McSaveney, M.J., 1986. Sedimentation and river containment on Waitangitaona alluvial fan—South Westland, New Zealand. Zeitschrift fur Geomorphologie 30, 215–230.
- Hayward, B.W., Grenfell, H.R., Reid, C.M., Hayward, K.A., 1999a. Recent New Zealand shallow-water benthic foraminifera: taxonomy, ecologic distribution, biogeography, and use in paleoenvironmental assessment. Institute of Geological and Nuclear Sciences Monograph 21.
- Hayward, B.W., Grenfell, H.R., Scott, D.B., 1999b. Tidal range of marsh foraminifera for determining former sea-level heights in New Zealand. New Zealand Journal of Geology and Geophysics 42, 395-413.
- Hayward, B.W., Scott, G.H., Grenfell, H.R., Carter, R., Lipps, J.H., 2004. Estimation of tidal elevation and salinity histories of sheltered harbours and estuaries using benthic foraminifera. The Holocene 14, 218-232.
- Hendey, N.I., 1964. An introductory account of the smaller algae of British coastal waters. Part V: Bacillariophyceae (Diatoms). Fisheries Investigation Series I. HMSO, London.
- Hull, A.G., 1986. Pre-A.D. 1931 tectonic subsidence of Ahuriri Lagoon, Napier, Hawke's Bay, New Zealand. New Zealand Journal of Geology and Geophysics 29, 75-82.
- Hustedt, F., 1927–1966. Die Kieselalagen Deutschlands, Osterreichs und der Schweiz (I–III). In: Rabenhorst, L. (Ed.), Kryptogamen-Flora von Deutschland, Osterreich under der Schweiz. 7. Akademische Verlagsgesellshaft, Leipzig.
- Hustedt, F., 1930. Bacillariophyta (Diatomaceae). In: Pascher, A. (Ed.), Die Susswasser-flora Mitteleuropas. Heft 10. Gustav Fischer Verlag, Jena.
- Hustedt, F., 1957. Die Diatomeenflora des Fluss-Systems der Weser im Gebiet der Hansestadt Bremen. Abhandlungen Naturwissenschaftlicher Verein, Bremen 34.
- John, J., 1983. The diatom flora of the Swan River estuary Western Australia. Bibliotecha Phycologica 64.
- Krammer, K, Lange-Bertalot, H., 1986–1991. Bacillariophyceae.

 Teil: Naviculaceae (1986), pp. 1–876; 2. Teil: Bacillariaceae,
 Epithemiaceae, Surirellaceae (1988), pp. 1–596; 3. Teil: Centrales,

881 882

883

884

885

886

887

888

889

890

891

892

893 894

895

896

897

898

899

900

901

902 903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

- Fragilariaceae, Eunotiaceae (1991), pp. 1–576; 4. Teil: Achnantha ceae, Kritische Erganzungen zu Navicula (Lineolatae) und Gom phonema Gesamtliteraturverzeichnis (1991), pp. 1–437. In: Pascher,
 A. (ed) Süswasserflora von Mitteleuropa, 2 (1–4). Gustav Fischer
 Verlag, Stuttgart.
- López-Buendía, A.M., Bastida, J., Querol, X., Whateley, M.K.G.,
 1999. Geochemical data as indicators of paleosalinity in coastal
 organic-rich sediments. Chemical Geology 157, 235–254.
- Loring, D.H., 1991. Normalization of heavy-metal data from estu arine and coastal sediments. ICES Journal of Marine Science 48,
 101-115.
- MacPherson, J.M., 1981. Hydrology of Okarito Lagoon and the inferred effects of selective logging in Okarito Forest. New Zealand
 Journal of Marine and Freshwater Research 15, 25-39.
- 843 McFadgen, B.G., Manning, M.R., 1990. Calibrating New Zealand 844 radiocarbon dates of marine shells. Radiocarbon 32, 229–232.
- Minoura, K., Nakaya, S., Uchida, M., 1994. Tsunami deposits in a
 lacustrine sequence of the Sanriku coast, Northeast Japan. Sedimentary Geology 89, 25-31.
- Minoura, K., Gusiakov, V.G., Kuratov, A., Takeuti, S., Svendsen, J.I.,
 Bondevik, S., Oda, T., 1996. Tsunami sedimentation associated
 with the 1923 Kamchatka earthquake. Sedimentary Geology 106,
 145–154.
- Molinaroli, E., De Falco, G., Rabitti, S., Portaro, R.A., 2000. Streamscanning laser system, electric sensing counter and settling grain analysis: a comparison using reference materials and marine sediments. Sedimentary Geology 130, 269–281.
- New Zealand Forest Service, 1984. Survey of the saltwater historic
 features. Unpublished New Zealand Forest Service Report, Hoki tika. 54 pp.
- Norris, R.J., Cooper, A.F., 1997. Erosional control on the structural
 evolution of range front faulting on the Alpine fault. Geological
 Society of New Zealand Miscellaneous Publication 95A.
- 862 Pascoe, J.D. (Ed.), 1952. The Great Journey: An Expedition to Explore
 863 the Interior of the Middle Island, New Zealand, 1846-1848.
 864 Pegasus Press, Christchurch. 111 pp.
- Round, F.E., Crawford, R.M., Mann, D.G., 1990. The Diatoms: Biology and Morphology of the Genera. Cambridge University Press,
 Cambridge.
- Scott, D.B., Medioli, F.S., 1986. Foraminifera as sea-level. in: van de
 Plassche, O. (Ed.), Sea-level Research: A Manual for the Collection and Evaluation of Data. Geo Books, Norwich.
- Shennan, I., Long, A.J., Rutherford, M.M., Green, F.M., Innes, J.B.,
 Lloyd, J.M., Zong, Y., Walker, K.J., 1996. Tidal marsh stratigraphy, sea-level change and large earthquakes, I: a 5000 year
 record in Washington, U.S.A. Quaternary Science Reviews 15,
 1023-1059.
- 876 Simonsen, R., 1967. Diatoms as indicators in the estuarine environment. Veroffentl. Inst. Meersforsch, Bremerhaven 11, 287–291.
- 878 Stuiver, M., Reimer, P.J., Bard, E., Beck, J.W., Burr, G.S., Hughen, 879 K.A., Kromer, B., McCormac, F.G., Plicht, J.V.D., Spurk, M.,

929

- 1998. INTCAL98 Radiocarbon Age Calibration, 24,000-0 cal BP. Radiocarbon 40, 1041–1083.
- Thomas, E., Varekamp, J.C., 1991. Paleo-environmental analyses of marsh sequences (Clinton, Connecticut): evidence for punctuated rise in relative sealevel during the latest Holocene. Journal of Coastal Research 11, 125–158.
- Van de Plassche, O., van der Borg, K., de Jong, A.F.M., 1998. Sea level climate correlation during the past 1400 yr. Geology 26, 319-322.
- Van der Werff, A., Huls, H., 1957-1974. Diatomeënflora van Nederland, vol. 1-10. Otto Koeltz Science Publishers, Koenigstein. Reprinted 1976.
- Vos, P.C., de Wolf, H., 1993a. Diatoms as a tool for reconstructing sedimentary environments in coastal wetlands; methodological aspects. Hydrobiologia 269/270, 285-296.
- Vos, P.C., de Wolf, H., 1993b. Reconstruction of sedimentary environments in Holocene coastal deposits of the southwest Netherlands; the Poortvliet boring, as a case study of palaeoenvironmental diatom research. Hydrobiologia 269/270, 297-306.
- Vos, P.C., de Wolf, H., 1994. Palaeoenvironmental research on diatoms in Early and Middle Holocene deposits in central and north Holland (The Netherlands). Netherlands Journal of Aquatic Ecology 28, 97-115.
- Wells, A., Goff, J., 2006. Coastal dune ridge systems as chronological markers of palaeoseismic activity; a 650-yr record from southwest New Zealand. The Holocene 16, 543-550.
- Wells, A.M., Stewart, G.H., Duncan, R.P., 1998. Evidence of widespread, synchronous, disturbance-initiated forest establishment in Westland, New Zealand. Journal of the Royal Society of New Zealand 28, 333-345.
- Wells, A.M., Yetton, M., Duncan, R.P., Stewart, G.H., 1999. Prehistoric dates of the most recent Alpine fault earthquakes, New Zealand. Geology 27, 995-998.
- Wells, A.M., Duncan, R.P., Stewart, G.H., 2001. Forest dynamics in Westland, New Zealand: the importance of large, infrequent earth-quake-induced disturbance. Journal of Ecology 89, 1006-1018.
- Wood, E.J.F., 1961. Studies on Australian and New Zealand diatoms, IV and V. Transactions of the Royal Society of New Zealand 88, 669-712.
- Wood, E.J.F., 1963. Studies on Australian and New Zealand diatoms, VI: tropical and subtropical species. Transactions of the Royal Society of New Zealand Botany 2, 189-218.
- Wood, E.J.F, Crosby, L.H., Cassie, V., 1959. Studies on Australian and New Zealand diatoms, III: descriptions of further discoid species. Transactions of the Royal Society of New Zealand 87, 211–219.
- Yetton, M., Wells, A., Traylen, N.J., 1998. The probability and consequences of the next Alpine fault earthquake. EQC Research Report, vol. 95/193. Wellington.

